

LIBRARY OF THE
UNIVERSITY OF ILLINOIS
AT URBANA-CHAMPAIGN

590.5

FI

v. 39

cop. 3



NATURAL HISTORY,
SURVEY

1871

1872

1873

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

FIELDIANA • ZOOLOGY

Published by

CHICAGO NATURAL HISTORY MUSEUM

Volume 39

FEBRUARY 27, 1959

No. 25

THREE NEW BIRDS FROM WEST AFRICA

AUSTIN L. RAND

CHIEF CURATOR, DEPARTMENT OF ZOOLOGY

AND

MELVIN A. TRAYLOR, JR.

ASSOCIATE CURATOR, DIVISION OF BIRDS

The new subspecies described below were encountered during the study of a collection of birds from Gabon, to be reported on elsewhere. For the loan of comparative material we are indebted to the authorities of the following institutions: Dean Amadon and Charles Vaurie, American Museum of Natural History; Kenneth Parkes, Carnegie Museum; James Greenway, Museum of Comparative Zoology.

Polipicus elliotii

Recently, *Polipicus johnstoni* of Mount Cameroon, with its races *schultzei* of Fernando Po and *kupeensis* of Mount Kupe, has been united with *P. elliotii* by Serle (Bull. Brit. Orn. Club, 72, 1952, p. 105), and in this action he is quite correct. *Polipicus elliotii* (sens. str.) has been considered monotypic, ranging from the Cameroon lowlands south to northern Angola and the Kasai, and east to the Ituri and Uganda. However, within *P. elliotii* there is a great deal of geographic variation, and at least one race must be separated.

Birds from the Cameroon agree most closely with Cassin's original description (Proc. Acad. Nat. Sci., 1863, p. 197). The type came from the Muni River, now the boundary between Spanish Guinea and Gabon. The diagnostic characters of the Cameroon population are the golden wash over much of the greenish upper parts, particularly the inner secondaries, coverts and scapulars; the heavy streaking below, extending onto the abdomen; and the dark yellow-olive wash over the breast and abdomen, the wash being darkest on the breast.

Library of Congress Catalog Card Number: 59-9479

Cassin particularly mentions the golden wash on the inner secondaries and the yellowish wash below being heaviest on the breast.

Gabon and Kasai specimens agree with Cameroon birds in the coloration and heavy streaking of the under parts but have the upper parts a clear yellow-green, with only a suggestion of golden wash on the secondaries and coverts.

East Congo-Uganda birds agree well with Gabon-Kasai specimens in the coloration of the upper parts and the width and extent of the ventral streaking. However, they lack the strong yellow-olive wash below that characterizes that population and the Cameroon specimens, having the ground color of the under parts a uniform pale yellow-green.

Angola specimens are at the opposite extreme from those of Cameroon, being the purest green above and having a pale yellow wash below, with the ventral streaking much reduced. The streaking on the breast is much narrower, and on the abdomen it is almost obsolete. This population is sufficiently distinct to be separated as:

***Polipicus elliotii gabela*, new subspecies**

Type.—Chicago Natural History Museum no. 224538, from 15 km. south of Gabela, Angola. Adult male, collected August 15, 1954, by Gerd Heinrich.

Diagnosis.—Compared to nominate *P. e. elliotii* of southern Cameroon, *gabela* has the upper parts a much clearer green, with very little golden wash; the dark streaking of the under parts is much reduced, being almost wanting on the abdomen; and the ground color of the under parts is pale yellowish, instead of yellow olive. *P. e. gabela* most nearly resembles *kupeensis* of British Cameroons, from which it is distinguished by the more buffy, less olive, ear coverts and sides of head, and the more whitish, less yellowish throat. There appears to be no significant variation in size within the mainland populations of the species.

Range.—Northwestern Angola.

Remarks.—The populations retained in *elliotii* are not uniform, but further division is not practicable. The Gabon and Congo populations resemble *gabela* in having more greenish upper parts, and upper Congo birds lack the strong olive wash below. All, however, are strongly streaked on the under parts and in this respect are readily separable from *gabela*.

The close resemblance between *gabala* and *kupeensis* is remarkable in view of their wide geographic separation; *kupeensis* is intermediate between the practically unstreaked *johnstoni* of Cameroon highlands and the heavily streaked *elliottii* of lowland Cameroons, and it was the discovery of this form that led Serle to unite the two species. The presence of a similar form at the opposite end of the range of *elliottii* strengthens this view.

The ranges and diagnostic characters of the recognizable races of *elliottii* are listed below:

schultzei.—Highlands of Fernando Po; smaller than *johnstoni* (not seen). Wing 85 mm. (Reichenow).

johnstoni.—Highlands of eastern Nigeria and British Cameroons (except Mount Kupe); greenish above, clear yellowish below, virtually unstreaked. Wing, 4 ♂, 91–93; 4 ♀, 86–91 mm.

kupeensis.—Mount Kupe, British Cameroons; like *johnstoni*, but moderately streaked below. Wing, 1 ♂, 92 mm.

elliottii.—Lowlands of British Cameroons east through southern Cameroon and the upper Congo to Uganda, and south through Gabon to the Kasai; always heavily streaked below, upper parts varying from golden green to green, lower parts from olive yellow to pale dull greenish-yellow. Wing, 13 ♂, 88–97 (av. 94.0); 10 ♀, 90–95 (av. 92.6 mm.).

gabala.—Northwestern Angola; green above and pale yellow below, only moderately streaked. Wing, 2 ♂, 93, 93; 1 ♀, 94 mm.

Specimens examined.—*johnstoni*: Cameroon Mountain, 4 ♂, 4 ♀. *kupeensis*: Mount Kupe, 1 ♂. *elliottii*: southern Cameroons, 9 ♂, 6 ♀; Gabon, M'Bigou, 1 ♀; Belgian Congo, Kasai, 1 ♂, Ituri, 2 ♂, 1 ♀, Rugege Forest, southeast of Lake Kivu, 1 ♂; Uganda, 2 ♀. *gabala*: Angola, Gabala, 1 ♂ (type), 2 ♀, Canzele, 30 km. west of Camabatela, 1 ♂.

Nectarinia superba

This species ranges throughout the Upper and Lower Guinea forests from Sierra Leone to Uganda and south to northern Angola. There is a cline of decreasing size from Uganda (wing of males 77–81) to Upper Guinea (wing of males 67–74), and a secondary north-south cline, Angola birds being almost as large as those from Uganda (wing of males 76–80). Variation in color, however, is not clinal. Uganda birds (*buvoma*) have the most extensive iridescent caps; those from Upper Guinea (*ashantiensis*) the least. Gabon and Angolan birds (*superba*) and those from Upper Guinea have the duller red under parts, while those from Nigeria and Cameroon have the brightest. There is no name available for this brightly colored population, so we propose to name it:

Nectarinia superba nigeriae, new subspecies

Type.—Chicago Natural History Museum no. 95910, from Ifon, Province of Ondo, Nigeria. Adult male collected by Rudyerd and Laura Boulton, August 10, 1934.

Diagnosis.—Male distinguished from all other races in being brighter, clearer red on the under parts. Iridescent cap as in nominate *superba*, of Lower Guinea; more extensive than in *ashantiensis* of Upper Guinea. Size intermediate between *superba* and the smaller *ashantiensis*, but nearer the former. Female not seen.

Measurements.—Type: wing 74, culmen from base 36. Five males: wing 72–76 (74.0); culmen 32–36 (34.5).

Range.—Southern Nigeria, intergrading through southern Cameroon with *superba*.

Remarks.—This new race is more closely allied to *superba* than to *ashantiensis*. The population of southern Cameroon is extremely variable. The under parts of individuals from the same localities vary from almost as bright as *nigeriae* to as dark as typical *superba* from Gabon and Angola. Populations from the last two localities are quite constantly dark dull red below. In the color of the under parts the males of *buvoma* from Uganda are intermediate between *nigeriae* and *superba*.

Measurements of adult males of the various populations examined are:

	Number of specimens	Wing	Culmen from base
<i>ashantiensis</i>			
Liberia	(2)	67, 70	31, 32
<i>nigeriae</i>			
Nigeria	(5)	72–76 (74.0)	32–36 (34.5)
(intermediate)			
Cameroon	(34)	73–79 (75.5)	33–39 (36.1)
<i>superba</i>			
Gabon	(5)	73–77 (75.2)	35–38 (36.4)
Angola	(10)	76–80 (77.4)	34–38 (36.0)
<i>buvoma</i>			
Uganda	(11)	77–81 (78.5)	32.5–38.5 (35.6)

Serinus capistratus

In the Heinrich collection of Angolan birds in Chicago Natural History Museum there is a fine series of four males, three females,

and five immature *Serinus capistratus* from the evergreen forests of Mount Soque and Mount Moco. When compared to males from adjoining regions in Angola, and from Gabon and the Kasai, the males are a much duller yellow on the under parts, and must be separated as:

***Serinus capistratus hildegardae*, new subspecies**

Type.—Chicago Natural History Museum no. 225994, from Mount Soque, 42 km. west-southwest of Luimbale, Huambo, Angola. Adult male, collected by G. Heinrich, August 28, 1954.

Diagnosis.—Males: similar to *capistratus*, but forehead, superciliaries and under parts duller, slightly olive yellow, rather than clear yellow. Females: darker, more olive yellow on the under parts, and as a consequence the dark streaking on the breast less conspicuous. Average somewhat larger than *capistratus*.

Measurements.—Wing: 4 males 65, 66, 70, 70; 3 females 65, 66, 67 mm.

Range.—Apparently confined to the evergreen forests on Mount Soque and Mount Moco, Huambo, Angola.

Specimens examined.—*capistratus*: Gabon, 3 ♂, 1 ♀; Angola, Canzele, 1 ♂, 1 ♀, Golungo Alto, 2 ♂ (one AMNH), Chitau, 1 ♂ (CM), Pungo Andongo, 1 ♂, 2 ♀ (AMNH), Canhoca, 1 ♂ (AMNH), Calamunjamba, 1 ♂ (AMNH). *hildegardae*: Mount Moco, 1 ♂, 1 ♀; Mount Soque, 3 ♂, 2 ♀, 2 im. ♂, 3 im. ♀.

Remarks.—Males of *capistratus* from northern Angola, Gabon and the Kasai are consistent in being a clear golden yellow on the under parts, and *hildegardae* is evidently confined to the mountain forests of Huambo. Wing measurements for *capistratus*: 11 males 61–67 (63.3); 4 females 61, 64, 64, 65. While not great the size difference is consistent.

There are five immature birds from Mount Soque, two of them beginning their molt into adult plumage. The immatures are patterned like the females, but are washed with dull olive green below. Even in this plumage the males are somewhat brighter than the females, though with just as much striping on the breast.

This subspecies is named in honor of Mrs. Gerd (Hildegard) Heinrich, who accompanied her husband in Angola and took an active part in the work of the trip.



UNIVERSITY OF ILLINOIS-URBANA



3 0112 027924254